

TITLE: Nucleoside syntheses. 19. C-Substitution of nucleosides with the aid of the Eschenmoser sulfide contraction
 AUTHOR(S): Vorbrueggen, Helmut; Krolikiewicz, Konrad
 CORPORATE SOURCE: Forschungslab., Schering A.-G., Berlin, Fed. Rep. Ger.
 SOURCE: Angewandte Chemie (1976), 88(21), 724-5
 CODEN: ANCEAD; ISSN: 0044-8249
 DOCUMENT TYPE: Journal
 LANGUAGE: German
 GI For diagram(s), see printed CA Issue.
 AB Treatment of thiopurine nucleosides I [R = SCH₂R₃ (R₃ = Bz, Me₃CO₂C, 4-O₂NC₆H₄CH₂); R₁ = H, Me₃SiNH; R₂ = Ac, Me₃Si] with strong base and Ph₃P gave C-alkyl nucleosides I (R = CH₂R₃, R₁ = H, NH₂) in 72-80% yields. Similarly prepared were II (X = CH, N; R = CH:C(OH)Ph, R₂ = H) and III (R = CH:C(OH)Ph, R₂ = H) from the corresponding II and III (R = SCH₂Bz, R₂ = Bz).
 IT 60363-87-3
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (Eschenmoser sulfide contraction of)
 RN 60363-87-3 ZCAPLUS
 CN Guanosine, 6-S-(2-oxo-2-phenylethyl)-6-thio-N-(trimethylsilyl)-2',3',5'-tris-O-(trimethylsilyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

